

POST-OP INSTRUCTIONS: CROWNS & BRIDGES

- Crowns and bridges usually take 2 or 3 appointments to complete. On the first appointment, the tooth/teeth are prepared, impressions are taken, and a temporary crown is placed on your tooth/teeth
- You may experience sensitivity, gum soreness, and slight discomfort on the tooth/teeth; it should subside after the placement of the permanent crown(s)
- For mild discomfort, alternate taking tylenol or ibuprofen **every 3-4 hours** or use a combination of both (as directed)
- When anesthesia is used, avoid chewing on your teeth until the numbness has worn off
- A temporary crown is usually made of plastic-based material. It can break if too much pressure is placed on it. The crown may also come off; if it does, save the crown and call our office. The temporary crown is placed to protect the tooth and prevent other teeth from moving. If it comes off it should be replaced as soon as possible. To avoid losing your temporary, do not chew on sticky or hard food (chewing gum, ice). Try to chew on the opposite side of the temporary as much as possible
- Continue your normal brushing but be careful while flossing around the temoriaries (remove the floss gently from the side). If it is difficult to get the floss between the temporary and surrounding teeth, refrain from flossing until you receive your permanent crown.
- After the permanent restoration is placed you may feel slight pressure for a few days. Also, the bite may feel different for a day or two. But if after 2-3 days the bite still feels uneven or if you feel discomfort when chewing on the tooth, call our office. Delaying the necessary adjustments may damage the tooth permanently.
- Call our office if you continue to be in pain

We are here to help if you need us! Please contact our office at 610-667-2770 or info@drweinstock.com if you have any questions or issues with these instructions.